

The Boltzmann Equation in Classical and Quantum Field Theory

Sangyong Jeon^{†,*}

[†]*Physics Department, McGill University,*

3600 University Street, Montreal, QC H3A-2T8, Canada and

**RIKEN-BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973*

Abstract

Improving upon the previous treatment by Mueller and Son, we derive the Boltzmann equation that results from a classical scalar field theory. This is obtained by starting from the corresponding quantum field theory and taking the classical limit with particular emphasis on the path integral and perturbation theory. A previously overlooked Van-Vleck determinant is shown to control the tadpole type of self-energy that can still appear in the classical perturbation theory. Further comments on the validity of the approximations and possible applications are also given.

I. INTRODUCTION

There are many situations in nature where classical field theory is the most direct and efficient way of describing the system under study. This happens when the occupation number becomes large enough that the quantum commutator becomes irrelevant. Such situations include most of macroscopic electro-magnetic systems and the dense systems created in colliding heavy ions. In studying heavy ion collisions, both the particle degrees of freedom and the classical field degrees of freedom becomes relevant: The final state of a collision consists mostly of the particle degrees of freedom whereas the initial state of two approaching nuclei can be efficiently described by a classical non-Abelian gauge field [1, 2, 3]. A natural framework to encompass both elements is the coupled system of classical field equation and the Boltzmann type of kinetic equation with the mean field[4].

An important problem in this framework is the conversion of classical field degrees of freedom into particle degrees of freedom and their subsequent thermalization. In this respect, the equivalence of the classical field theory and the Boltzmann equation proposed by Mueller and Son[5] is significant since it has a potential of providing a consistent framework for the thermalization in heavy ion collisions. The current paper improves on the work of Mueller and Son in the following aspects.

First, going from quantum to classical many body theory almost always involves the Wigner function [4, 6, 7]. In Ref.[5], this point was overlooked. In this work, we show that the Wigner functional does appear in the formulation due to the fact that the density operator is in general non-diagonal.

Second, the Feynman rules used in Ref.[5] for the classical field are a mixture of quantum and classical ones: The vertex rule is classical but the propagators are quantum mechanical. We show that consistent formulation only involves classical vertices and propagators.

Third, formulation of classical path integral involves a functional determinant related to the Jacobi field. Therefore, perturbation theory based on classical path integral involves ghost fermions [8, 9]. It turned out that the classical perturbation theory as formulated in Ref.[5] still contains a quantum effect in the form of the tadpole self-energy. The ghost contribution is shown to compensate this remaining quantum effect, thus making the theory consistently classical.

Despite these gaps in the formulation, the final result in Ref.[5] is essentially correct in

the $f \gg 1$ limit. The goal of this paper is to provide a more consistent derivation of the kinetic equation corresponding to the classical scalar field starting from the quantum scalar field. We take particular care in treating the path integral and the perturbation theory in a consistent manner. We also comment on the validity of this classical method and possible use in the context of understanding the thermalization in heavy ion collisions.

II. CLASSICAL PATH INTEGRAL FROM QUANTUM PATH INTEGRAL

To be specific, consider a real scalar field theory defined by the following Hamiltonian

$$H = \int d^3x \left(\frac{\pi^2}{2} + \frac{(\nabla\phi)^2}{2} + V(\phi) \right) \quad (1)$$

The equations of motion are

$$\dot{\phi} = \frac{\delta H}{\delta \pi} = \pi \quad (2)$$

$$\dot{\pi} = -\frac{\delta H}{\delta \phi} \equiv E(\phi) \quad (3)$$

where we defined $E(\phi) = \nabla^2\phi - V'(\phi)$ for notational convenience.

In many publications since 1980's E. Gozzi and his collaborators have been extensively studying the properties of classical path integrals (See Refs.[8, 9] and references therein.). The starting point in these studies is the following transition probability between two points in the phase space

$$P[\varphi_f, \pi_f^{\varphi}; t_f | \varphi_i, \pi_i^{\varphi}; t_i] = \delta[\varphi_f - \varphi_c(t_f | \varphi_i, \pi_i^{\varphi})] \delta[\pi_f^{\varphi} - \pi_c^{\varphi}(t_f | \varphi_i, \pi_i^{\varphi})] \quad (4)$$

Here φ and π^{φ} are the generalized coordinate and momentum and φ_c and π_c^{φ} are the solutions of Eqs. (2) and (3) with the boundary conditions $\varphi_c(t_i | \varphi_i, \pi_i^{\varphi}) = \varphi_i$ and $\pi_c^{\varphi}(t_i | \varphi_i, \pi_i^{\varphi}) = \pi_i^{\varphi}$. The evolution of the initial phase space density $\rho_{\text{cl}}[\varphi, \pi^{\varphi}; t_i]$ is given by

$$\rho_{\text{cl}}[\varphi, \pi^{\varphi}; t] = \int [d\varphi_i][d\pi_i^{\varphi}] P[\varphi, \pi^{\varphi}; t | \varphi_i, \pi_i^{\varphi}; t_i] \rho_{\text{cl}}[\varphi_i, \pi_i^{\varphi}; t_i] \quad (5)$$

In this paper we use the square bracket notation $\int [d\phi]$ to indicate a functional integral at a fixed time and the curly D notation $\int \mathcal{D}\phi = \int \prod_{i=1}^N [d\phi_i]$ to indicate a functional integral over space and time. We also implicitly absorb any normalization constants into the definition of $[d\phi]$.

To turn Eq.(5) into a path integral, we use the fact that transition probabilities satisfy

$$\int dy P[z|y]P[y|x] = P[z|x] \quad (6)$$

Dividing the time between the final time $t_f = t_{N+1}$ and the initial time $t_i = t_0$ into $N+1$ equal intervals we obtain

$$P[\phi_{N+1}, \pi_{N+1} | \phi_0, \pi_0] = \int \prod_{i=1}^N [d\phi_i][d\pi_i] \prod_{j=0}^N P[\phi_{j+1}, \pi_{j+1} | \phi_j, \pi_j] \quad (7)$$

where we have suppressed the time arguments in P .

For sufficiently small $\Delta t = (t_f - t_i)/(N+1)$, we should be able to solve the classical equations of motion approximately. The simplest finite difference method is the Euler method based on the following approximation of the time derivative

$$\dot{f}(t_k) \approx \frac{f(t_k) - f(t_{k-1})}{\Delta t} + O(\Delta t^2) \quad (\text{Backward Euler}) \quad (8)$$

or

$$\dot{f}(t_k) \approx \frac{f(t_{k+1}) - f(t_k)}{\Delta t} + O(\Delta t^2) \quad (\text{Forward Euler}) \quad (9)$$

We use here the Backward Euler method for φ

$$\varphi_{k+1} = \varphi_k + \pi_{k+1}^\varphi \Delta t + O(\Delta t^2) \quad (10)$$

and the Forward Euler method for π^φ

$$\pi_{k+1}^\varphi = \pi_k^\varphi + E(\varphi_k) \Delta t + O(\Delta t^2) \quad (11)$$

so that φ_{k+1} can be expressed solely in terms of φ_k and π_k^φ . This is, of course, not the only choice. Different discretization method in general leads to different form of the discretized path integral although they are all equivalent in the $\Delta t \rightarrow 0$ limit.

For small enough Δt then,

$$P[\varphi_{j+1}, \pi_{j+1}^\varphi | \varphi_j, \pi_j^\varphi] = \delta[\varphi_{j+1} - \varphi_j - \pi_{j+1}^\varphi \Delta t] \delta[\pi_{j+1}^\varphi - \pi_j^\varphi - E(\varphi_j) \Delta t] \quad (12)$$

With this form, it is easy to check that the preservation of probability

$$\int [d\varphi_{N+1}][d\pi_{N+1}^\varphi] \rho_{\text{cl}}[\varphi_{N+1}, \pi_{N+1}^\varphi; t_{N+1}] = \int [d\varphi_0][d\pi_0^\varphi] \rho_{\text{cl}}[\varphi_0, \pi_0^\varphi; t_0] \quad (13)$$

is trivially satisfied. Using dummy variables χ and π^χ to express the δ -functionals, we finally get the classical path integral for the evolution of the density functional

$$\begin{aligned} \rho_{\text{cl}}[\varphi_{N+1}, \pi_{N+1}^\varphi; t_{N+1}] &= \int \prod_{j=0}^N [d\varphi_j] \prod_{k=1}^{N+1} [d\pi_k^\chi] \exp \left(i \sum_{j=0}^N \pi_{j+1}^\chi (\varphi_{j+1} - \varphi_j - \pi_{j+1}^\varphi \Delta t) \right) \\ &\times \int \prod_{l=0}^N [d\pi_l^\varphi] \prod_{k=0}^N [d\chi_k] \exp \left(-i \sum_{k=0}^N \chi_k (\pi_{k+1}^\varphi - \pi_k^\varphi - E(\varphi_k) \Delta t) \right) \\ &\times \rho_{\text{cl}}[\varphi_0, \pi_0^\varphi; t_0] \end{aligned} \quad (14)$$

How closely can we reproduce the classical path integral Eq.(14) from the corresponding quantum path integral? To answer this question, we need to go back to the basics of deriving a quantum mechanical path integral. Given a Hamiltonian operator $\hat{H}$, we would like to rewrite the matrix element of the evolution operator, $\langle \phi_f | \hat{U}(t_f, t_i) | \phi_i \rangle$, as a path integral. Dividing the time interval into many small segments and using the fact that $\hat{U}(t, t') \hat{U}(t', t'') = \hat{U}(t, t'')$, we can write

$$\langle \phi_{N+1} | \hat{U}(t_{N+1}, t_0) | \phi_0 \rangle = \langle \phi_{N+1} | \hat{U}(t_{N+1}, t_N) \hat{U}(t_N, t_{N-1}) \cdots \hat{U}(t_1, t_0) | \phi_0 \rangle \quad (15)$$

There are many ways of inserting the resolutions of identity to turn this expression into a path integral just as there are many ways of discretizing the classical equation of motion. The prescription which most closely resembles Eqs.(10) and (11) turned out to be inserting

$$1 = \int [d\phi_k] [d\pi_k] |\phi_k\rangle \langle \phi_k | \pi_k \rangle \langle \pi_k | \quad (16)$$

between $\hat{U}(t_{k+1}, t_k)$ and $\hat{U}(t_k, t_{k-1})$ for all $1 \leq k \leq N$ and inserting

$$1 = \int [d\pi_{N+1}] |\pi_{N+1}\rangle \langle \pi_{N+1} | \quad (17)$$

between $\langle \phi_{N+1} |$ and $\hat{U}(t_{N+1}, t_N)$. In this way, we get

$$\begin{aligned} \langle \phi_{N+1} | \hat{U}(t_{N+1}, t_0) | \phi_0 \rangle &= \int \prod_{k=1}^N [d\phi_k] \int \prod_{k=1}^{N+1} [d\pi_k] \\ &\times \exp \left(i \sum_{k=1}^{N+1} \pi_k (\phi_k - \phi_{k-1}) - i \sum_{k=1}^{N+1} H(\pi_k, \phi_{k-1}) h \right) \end{aligned} \quad (18)$$

The time evolution of a density operator needs two such path integrals since the matrix element

$$\begin{aligned} \langle \phi_f | \hat{\rho}(t_f) | \tilde{\phi}_f \rangle &= \langle \phi_f | \hat{U}(t_f, t_i) \hat{\rho}(t_i) \hat{U}(t_i, t_f) | \tilde{\phi}_f \rangle \\ &= \int [d\phi_i] [d\tilde{\phi}_i] \langle \phi_f | \hat{U}(t_f, t_i) | \phi_i \rangle \langle \phi_i | \hat{\rho}(t_i) | \tilde{\phi}_i \rangle \langle \tilde{\phi}_i | \hat{U}(t_i, t_f) | \tilde{\phi}_f \rangle \end{aligned} \quad (19)$$

involves two matrix elements of the evolution operator $\hat{U}$. To bring out classical features more clearly, define

$$\varphi = (\phi + \tilde{\phi})/2 \quad (20)$$

$$\chi = \phi - \tilde{\phi} \quad (21)$$

where ϕ belongs to the path integral representation of $\langle \phi_f | \hat{U}(t_f, t_i) | \phi_i \rangle$ and $\tilde{\phi}$ belongs to the path integral representation of $\langle \tilde{\phi}_i | \hat{U}(t_i, t_f) | \tilde{\phi}_f \rangle$.

Introducing the Wigner functional $\rho_W(\varphi, \pi^\varphi)$ as

$$\langle \varphi + \chi/2 | \hat{\rho} | \varphi - \chi/2 \rangle = \int [d\pi^\varphi] \rho_W[\varphi, \pi^\varphi] e^{i\chi\pi^\varphi} \quad (22)$$

we get

$$\begin{aligned} \rho_W[\varphi_{N+1}, \pi_{N+1}^\varphi] &= \int [d\varphi_0][d\chi_0] \rho_W[\varphi_0, \pi_0^\varphi] \\ &\times \int \prod_{k=1}^N [d\varphi_k] \int \prod_{k=1}^N [d\chi_k] \int \prod_{k=0}^N [d\pi_k^\varphi] \int \prod_{k=1}^{N+1} [d\pi_k^\chi] \\ &\times \exp \left(i \sum_{k=0}^N \pi_{k+1}^\chi (\varphi_{k+1} - \varphi_k - \pi_{k+1}^\varphi \Delta t) \right) \\ &\times \exp \left(-i \sum_{k=0}^N \chi_k (\pi_{k+1}^\varphi - \pi_k^\varphi - E(\varphi_k) \Delta t) + O(\chi^3) \right) \end{aligned} \quad (23)$$

Note that without the Wigner transformation of the initial and the final distributions, the exponents cannot be arranged in the above finite-difference form that includes the end points.

The difference between Eq.(23) and the classical expression Eq.(14) is the $O(\chi^3)$ term inside the exponential. (There are no terms even in χ in this expression because the exponent must change sign under the exchange $\phi \leftrightarrow \tilde{\phi}$ or equivalently, $\chi \rightarrow -\chi$.) If we drop the $O(\chi^3)$ terms, then we have exactly the same *evolution kernel* as the classical theory. However, even in this case, this does not mean that ρ_W will be truly classical. If the initial Wigner functional contains quantum information, then ρ_W at any later times will still contain it.

In perturbation theory, dropping $O(\chi^3)$ terms means ignoring any Feynman diagrams that may contain a loop or loops made up of the *off-shell* propagators. The validity of such operation will be shortly discussed in the next section where we discuss the Feynman rules.

To summarize this section, the omission of the terms non-linear in χ results in the classical *evolution* of the initial distribution. The crucial difference between our analysis here and that of Ref.[5] is the appearance of the Wigner functional which depends both on the

coordinate (φ) and the momentum (π^φ). This is important in the perturbation theory since the propagators strongly depend on the initial distribution. We therefore discuss the propagators and the Feynman rules next.

III. PROPAGATORS AND FEYNMAN RULES

Having obtained the *classical* path integral from a quantum one, we can now derive Feynman rules for the classical perturbation theory following the same procedure as in the quantum case. To do so, consider the generating functional with the source terms $J_\varphi\varphi - J_\chi\chi$ and the restrictions that the source terms vanish at the end points t_i and t_f . Carrying out π^χ integrals and χ_0 integral results in

$$\delta \left[\frac{\varphi_1 - \varphi_0}{h} - \pi_0^\varphi - E(\varphi_0)h \right] \prod_{k=0}^N \delta \left[\varphi_{k+1} - \varphi_k - \pi_{k+1}^\varphi h \right]$$

We can then carry out all π_k^φ integrals to get

$$\begin{aligned} \mathcal{Z}[J_\varphi, J_\chi] &= \int \prod_{k=0}^{N+1} [d\varphi_k] \int \prod_{k=1}^N [d\chi_k] \rho_W[\varphi_0, (\varphi_1 - \varphi_0)/h - E(\varphi_0)h] \\ &\quad \times \exp \left(-i \sum_{k=1}^N \chi_k \left(\frac{\varphi_{k+1} - 2\varphi_k + \varphi_{k-1}}{h} - E(\varphi_k)h + J_k^\chi h \right) + i \sum_{k=1}^N J_k^\varphi \varphi_k h \right) \\ &= \int \mathcal{D}\varphi \mathcal{D}\chi \rho_W[\varphi_i, \dot{\varphi}_i] \\ &\quad \times \exp \left(i \int \chi \left(-\partial^2 \varphi - m^2 \varphi - \frac{\lambda}{3!} \varphi^3 - J_\chi \right) + i \int J_\varphi \varphi \right) \end{aligned} \quad (24)$$

where we switched to the continuum notation for simplicity.

It is tempting now to develop a perturbation theory from this expression. But we are not done yet. Note that the initial distribution depends on the value of the field and its time derivative. But the integration is over the field values only. To treat $\rho_W[\varphi_i, \dot{\varphi}_i]$ as a proper weight we need to transform the measure to include integration over $\dot{\varphi}_i$. In classical field theory, this can be most easily done by noting that to solve a second order differential equation requires specifying either the values of the field and its first time derivative at the initial time or the values of the field at the initial and the final time. Therefore, barring caustic points, there is a one-to-one correspondence between the value of φ at t_f and the value of $\dot{\varphi}$ at t_i . The Jacobian of this transformation is

$$\mathcal{J} = \left| \frac{\delta \varphi_f}{\delta \dot{\varphi}_i} \right| = \left| \frac{\delta^2 S}{\delta \varphi_i \delta \varphi_f} \right|^{-1} \quad (25)$$

where S is the action integral. This Jacobian is related to the Van-Vleck determinant

$$\mathcal{J}[\varphi] = \left| \text{Det}(\partial^2 + m^2 + (\lambda/2)\varphi^2) \right| \quad (26)$$

Changing the variable φ_f to $\dot{\varphi}_i$ then gives

$$\begin{aligned} \mathcal{Z}[J_\varphi, J_\chi] &= \int [d\pi_i][d\varphi_i] \rho_W[\varphi_i, \pi_i] \int_{\varphi_i, \pi_i} \mathcal{D}\varphi \mathcal{D}\chi \\ &\times \mathcal{J}[\varphi] \exp \left[i \int \chi \left(-\partial^2 \varphi - m^2 \varphi - \frac{\lambda}{3!} \varphi^3 - J_\chi \right) + i \int J_\varphi \varphi \right] \end{aligned} \quad (27)$$

where we have renamed $\dot{\varphi}_i = \pi_i$ and there is no $[d\varphi_f][d\chi_f]$ integral. One can easily check that in the limit $J_\varphi \rightarrow 0$, $\mathcal{Z} \rightarrow 1$. This is as it should be since $\mathcal{Z}$ reduces to a trace of a density operator in this limit.

Note that without the Jacobian $\mathcal{J}$, the $J_\varphi \rightarrow 0$ limit will not in general yield 1. This is because performing χ integral produces $\delta[(\partial^2 \varphi + m^2 \varphi + (\lambda/3!)\varphi^3) + J_\chi]$ which is then integrated over φ to yield $1/\mathcal{J}$. The exact value of the Jacobian, however, depends on how one discretizes the equation of motion. This may sounds peculiar, but this phenomenon is not new. In stochastic dynamics, it is well known that this kind of functional Jacobian depends on the discretization prescription[10]. Furthermore, the Jacobian can be made constant if one choose a particular prescription. This was also noticed in Ref.[11]. Later in this section, we will comment more on the role of the determinant. For now we simply set it to a constant and ignore it. As shown shortly, this amounts to the following diagrammatic rule:

- (i) Omit diagrams containing tadpoles.

To get the rest of the Feynman rules, let us examine the two main ingredients of perturbation theory: Propagators and interaction vertices. The forms of the interaction vertices are fully determined by the interaction Lagrangian. In the case of the classical $\lambda\phi^4$ theory, the corresponding Feynman rule is:

- (ii) Assign $-i\lambda$ to each $\lambda\varphi^3\chi$ vertex.

To determine propagators, one must specify what the *un-perturbed state* is. In the quantum theory, this is the perturbative vacuum which can fluctuate into particle-antiparticle pairs. Therefore, the expectation value $\langle 0|\hat{\phi}(x)\hat{\phi}(y)|0\rangle$ is non-zero even if $\langle 0|\hat{\phi}(x)|0\rangle = 0$. On the

other hand, in the classical vacuum, both of these quantities are zero because the classical vacuum cannot fluctuate. It is literally a state where nothing exists.

Propagators are determined by the free field limit. Following the derivation in the last section, it is easy to see that without the self-interaction, the generating function quickly reduces to

$$\begin{aligned} \mathcal{Z}[J_\varphi, J_\chi] &= \int [d\pi_i][d\varphi_i] \rho_W[\varphi_i, \pi_i] \\ &\times \int_{\varphi_i, \pi_i} \mathcal{D}\varphi \mathcal{J}[\varphi] \delta[-\partial^2\varphi - m^2\varphi + J_\chi] \exp\left[i \int J_\varphi \varphi\right] \end{aligned} \quad (28)$$

regardless of whether it's the quantum case or the classical case. Yet for both cases, the combination $\varphi = (\phi + \tilde{\phi})/2$ satisfies the classical equation of motion whose solution is given by

$$\varphi = \left(\varphi_i \cos(E_k(t - t_i)) + \pi_i \frac{\sin(E_k(t - t_i))}{E_k} \right) + \int_{t_i}^t dt' G_{\text{ret}}(t - t') J_\chi(t') \quad (29)$$

The fact that φ satisfies the classical equation of motion even in the fully quantum case is somewhat unexpected. However, it shows explicitly that the form of the initial distribution functional ρ_W plays an essential role in distinguishing the quantum and the classical cases.

Let's consider the purely classical case first. In equilibrium, the density functional is given by the classical Boltzmann factor

$$\rho_{\text{cl}}[\varphi_i, \pi_i] = \frac{e^{-\beta H_{\text{cl}}}}{Z_{\text{cl}}} = \frac{1}{Z_{\text{cl}}} \exp\left(-\beta \sum_k (\pi_i(k)^2 + E_k^2 \varphi_i(k)^2)/2\right) \quad (30)$$

where $\pi_i(k)$ and $\varphi_i(k)$ are the Fourier components of π_i and φ_i respectively. Using the above classical solution and carrying out the φ_i and π_i integral then yield

$$\mathcal{Z}_{\text{cl}}[J_\varphi, J_\chi] = \exp\left[-\sum_k \frac{T}{2E_k^2} \int_{t_i}^{t_f} dt \int_{t_i}^{t_f} dt' J_\varphi(t) \cos(E_k(t - t')) J_\varphi(t') + i \int J_\varphi G_{\text{ret}} J_\chi\right] \quad (31)$$

where we have again suppressed the momentum indices for J_φ and J_χ . The resulting Feynman rules are:

(iii) In equilibrium, the propagators are

$$G_{\text{cl}}^{\varphi\varphi}(p) = (T/E_p) 2\pi \delta(p^2 - m^2) \quad (32)$$

$$G_{\text{cl}}^{\varphi\chi}(p) = \frac{i}{p^2 - m^2 + i\epsilon p_0} \quad (33)$$

$$G_{\text{cl}}^{\chi\varphi}(p) = \frac{i}{p^2 - m^2 - i\epsilon p_0} \quad (34)$$

$$G_{\text{cl}}^{\chi\chi}(p) = 0 \quad (35)$$

The same set of rules were also used in Refs.[12, 13, 14, 15] as the classical limit of the quantum rules.

The $\varphi\chi$ propagator is the response function which controls generation of classical field by a source. Hence it must be there be it classical or quantum (*c.f.* Eq.(29)). However, in the $T \rightarrow 0$ limit, $\varphi\varphi$ propagator vanishes due to the fact that classical vacuum does not fluctuate. Note also that the classical equilibrium density function is T/E_k just as in the case of electromagnetic waves in a cavity. Therefore, if one is to extend this formalism to an out-of-equilibrium situation, one should use

(iii') In non-equilibrium,

$$G_{\text{cl}}^{\varphi\varphi}(p, X) = f_{\text{cl}}(p, X) 2\pi \delta(p^2 - m^2) \quad (36)$$

where $f_{\text{cl}}(p, X)$ is the density function to be determined and $G_{\text{cl}}^{\varphi\varphi}(p, X)$ is the Wigner transformed 2 point function

$$G_W(p, X) = \int d^4r e^{ir^\mu p_\mu} G(x, y) \quad (37)$$

with $r = x - y$ and $X = (x + y)/2$.

The appearance of the on-shell δ -function in $G_{\text{cl}}^{\varphi\varphi}$ propagator deserves deeper consideration. Suppose that the initial $\rho[\varphi_i, \chi_i]$ is actually independent of χ_i . That is, $\hat{\rho}$ is diagonal in ϕ . In that case, the Wigner functional contains a δ functional $\delta[\pi_i]$. For an illustration, suppose

$$\rho'_W[\varphi_i, \pi_i] = \frac{1}{Z'} \exp \left(-\beta \sum_k E_k^2 \varphi_i(k)^2 / 2 \right) \delta[\pi_i] \quad (38)$$

In that case, carrying out φ_i and π_i^φ integrals results in

$$\mathcal{Z}'[J_\varphi, J_\chi] = \exp \left[- \sum_k \frac{T}{2E_k^2} \left(\int_{t_i}^{t_f} dt J_\varphi(t) \cos(E_k(t - t_i)) \right)^2 + i \int J_\varphi G_{\text{ret}} J_\chi \right] \quad (39)$$

Since there is no compensating sine term, the $\varphi\varphi$ Green function no longer exhibits time-translation invariance. The $\varphi\varphi$ correlator is given by

$$G^{\varphi\varphi}(t, t') = \frac{T}{E_k^2} \cos(E_k(t - t_i)) \cos(E_k(t' - t_i)) \quad (40)$$

Since this is not a function of $t - t'$, this form of the propagator *does not* conserve energy at each vertex. The usual momentum space Feynman diagram techniques will not work.

Our ability to use Feynman rules with the usual conservation 4-momentum at each vertex depends on whether we get a function only of $t - t'$ for the $\varphi\varphi$ correlator. In equilibrium, the propagators depend only on the difference of the two coordinates. Hence, the total 4-momentum is conserved at each vertex. In non-equilibrium situations, the propagators depend both on the difference $r = x - y$ and the sum $X = (x + y)/2$. Wigner transformation assigns ‘momentum’ to the difference as done in Eq.(37). However, unless the X dependence of $G_W(p, X)$ is slow, the momenta that enter each vertex are not conserved. For the derivation of the Boltzmann equation, this slow dependence is therefore essential.

The above Feynman rules are used to calculate the perturbative corrections in the classical field theory. Just as loops appear in the quantum Feynman diagrams, loops appear in the classical Feynman diagrams as well [5]. This looks like a violation of the theorem that states the classical field theory corresponds to the sum of all tree diagrams. However, this theorem is derived in the context of the vacuum theory. It does not necessarily apply to the in-medium case. Furthermore, the theorem is derived considering only the un-cut diagrams where as the in-medium theory necessarily involves cut diagrams (that include $\varphi\varphi$ propagators). Nevertheless, if this theorem is not to be violated, we must at least show that the vacuum loops do not appear in our classical theory.

That this is indeed the case can be shown as follows. A vacuum loop in a given diagram must not contain the density function f_{cl} . Since f_{cl} appears only in the $\varphi\varphi$ propagator, any potential vacuum loop diagram must consist entirely of $\chi\varphi$ and $\varphi\chi$ propagators. Now with only the $\lambda\chi\varphi^3$ vertex, χ is conserved in the sense that it cannot split into multiple χ ’s. Hence one can follow χ as if it is a fermion. What flows along this line is the time since the $\varphi\chi$ propagator is a retarded one and the $\chi\varphi$ propagator is an advanced one. Therefore, a loop made of $\varphi\chi$ and/or $\chi\varphi$ propagators implies existence of a time-loop which is impossible. It follows that there is no vacuum loop contribution in the classical theory if the loop involves more than a single propagator.

The only remaining possibility is a single loop formed by a $\varphi\chi$ propagator as shown in Fig.1 which corresponds to

$$\Sigma_{1-loop} = \frac{\lambda}{2} G_{ret}(t = 0) \quad (41)$$

The value of this expression is actually ambiguous as it involves $\theta(0)$. This is where the Van-Vleck determinant Eq.(26) plays its role. To determine the role of $\mathcal{J}$ in perturbation




FIG. 1: Tadpole diagram in the classical perturbation theory.

theory, we may exponentiate the determinant using Fadeev-Popov type of ghost fields c and $\bar{c}$

$$\text{Det}(\partial^2 + m^2 + (\lambda/2)\varphi^2) \rightarrow \int \mathcal{D}c \mathcal{D}\bar{c} \exp \left(i \int \bar{c}(\partial^2 + m^2 + (\lambda/2)\varphi^2)c \right) \quad (42)$$

Since there is no in-medium part for the ghost field, the corresponding propagators should be

$$G_Q^{c\bar{c}} = \frac{i}{p^2 - m^2 + i\epsilon p_0} \quad (43)$$

$$G_Q^{\bar{c}c} = \frac{i}{p^2 - m^2 - i\epsilon p_0} \quad (44)$$

with the rule that a ghost-loop gets an additional factor of (-1) .

First of all, ghost propagators must form a loop since no external line can be a ghost and there can be no ghost excitations in the medium. However, since the ghost propagator is a retarded one, a ghost loop made of 2 or more ghost lines must vanish. A loop made up of a single ghost line is then the only possible non-zero diagram

$$\Sigma_{1\text{-loop}}^{\text{ghost}} = -\frac{\lambda}{2} G_{\text{ret}}(t=0) \quad (45)$$

This exactly cancels the tadpole contribution in Eq.(41). Thus the ghost saves the classicality of the perturbation theory by removing the ambiguity of defining $\theta(0)$. If we define $\theta(0) = 0$, then the ghost field contribution disappears and the Van Vleck determinant can be set to a constant. Again, as mentioned earlier, this is a known phenomenon. In the study of stochastic dynamics, it is well known that how one discretizes can change the form of Van Vleck determinant even making it a constant[10]. In Ref.[5], since all loops that are formed with $\varphi\chi$ propagators and/or $\chi\varphi$ propagators are set to zero, one might say that $\theta(0)$ was implicitly defined to be zero in that paper.

To determine the validity of the above classical perturbation theory as a approximation of the underlying quantum theory, let us now consider the quantum case. The form of the free field Wigner function can be easily obtained from the ground state wavefunction of the simple harmonic oscillator in the Euclidean space (For instance, see Ref.[16].).

$$\rho_Q[\varphi_i, \pi_i] = \frac{1}{Z_Q} \exp \left(- \sum_k \frac{\tanh(E_k \beta / 2)}{E_k} \left(\pi_i(k)^2 + E_k^2 \varphi_i(k)^2 \right) \right) \quad (46)$$

In the small β or large $T = 1/\beta$ limit, this goes over to the classical case. Again using the classical solution and carrying out the φ_i and π_i integral yield

$$\mathcal{Z}_Q[J_\varphi, J_\chi] = \exp \left[- \sum_k \frac{\coth(E_k \beta / 2)}{4E_k} \int_{t_i}^{t_f} dt \int_{t_i}^{t_f} dt' J_\varphi(t) \cos(E_k(t - t')) J_\varphi(t') + i \int J_\varphi G_{\text{ret}} J_\chi \right] \quad (47)$$

The propagators are then given by

$$G_Q^{\varphi\varphi} = (1/2 + n_{\text{BE}}(E_p)) 2\pi \delta(p^2 - m^2) \quad (48)$$

$$G_Q^{\varphi\chi} = \frac{i}{p^2 - m^2 + i\epsilon p_0} \quad (49)$$

$$G_Q^{\chi\varphi} = \frac{i}{p^2 - m^2 - i\epsilon p_0} \quad (50)$$

$$G_Q^{\chi\chi} = 0 \quad (51)$$

where $n_{\text{BE}}(E_p) = 1/(e^{E_p/T} - 1)$ is the Bose-Einstein distribution. The corresponding non-equilibrium propagator is

$$G_Q^{\varphi\varphi} = (1/2 + f_Q) 2\pi \delta(p^2 - m^2) \quad (52)$$

where f_Q is now the density function to be determined. In Ref.[5], these quantum form of propagators are used with the classical field Feynman rules (i) and (ii) above. If $f_Q \gg 1/2$, there is very little difference between this set of quantum propagators and the set of classical propagators. However, one of the points made in Ref.[5] was that keeping the $1/2$ term in Eq.(52) makes the classical kinetic equation to match up with the quantum one up to the next-to-leading order in f . In view of the fact that this is actually mixing quantum and classical descriptions, it deserves a more in-depth study. We will shortly do so in the next section.

Since we have explicit forms of the quantum and the classical propagators now, we can discuss the validity of the classical perturbation expansion. The differences between the classical and the quantum Feynman rules are as follows (excluding the tadpole diagrams):

- (i) The quantum $\varphi\varphi$ propagator has $1/2 + f$ whereas the classical one has just f .
- (ii) The quantum interaction includes $\lambda\chi^3\varphi/24$ term whereas this is missing in the classical case.

The first of these items indicates that the classical approximation is valid in the large f limit, $f \gg 1$ so that the appearance of $1/2$ does not make a difference. But if f is too big,




FIG. 2: The first order vertex correction for the classical field.

then potentially perturbative corrections can be as large or larger than the leading order contribution. To be concrete, consider the vertex correction depicted in Fig. 2. Compared to the bare vertex, the correction term is smaller by a factor of $O(\lambda f_{\text{cl}})$. Therefore, the *classical* perturbation theory is valid when $f_{\text{cl}} \gg 1$ but also $\lambda f_{\text{cl}} \ll 1$. This is the same conclusion reached in Ref.[5] but in a heuristic way.

The next question we should ask is how big the size of the quantum correction is. To have an estimate, consider substituting a $\lambda\varphi^3\chi$ vertex with a $\lambda\chi^3\varphi$ vertex in a classical perturbation theory diagram. Since there is no $\chi\chi$ propagator, the only way this substitution is possible is when the inserted vertices are connected with three φ field and one χ field as depicted in Fig. 3. In the large f limit, the quantum contribution is $O(1/f^2)$ smaller than the purely classical contribution. Therefore at a given order in λ , the quantum correction is always $O(1/f^2)$ smaller than the classical one. Again, this is the same conclusion as in Ref.[5].

These estimates also indicate that there is a limit that we can trust the classical perturbation theory depending on the relative size of $1/f$ and λ . Suppose that $f = O(1/\sqrt{\lambda})$. In the small λ limit, this fulfills $f \gg 1$ as well as $\lambda f \ll 1$. However, since $1/f^2 = O(\lambda)$, only the leading order classical perturbative correction is reliable. The leading order (in λ) quantum correction in this case is as big as the second order classical correction.




FIG. 3: Substituting $\lambda\phi^3\chi$ with $\lambda\chi^3\phi$.

IV. COMPARISON OF QUANTUM AND CLASSICAL BOLTZMANN EQUATIONS

Having derived the Feynman rules, the derivation of Boltzmann equation can now proceed as in Ref.[5] for the $\lambda\phi^4$ theory. For convenience, here we show the results from Ref.[5]. The collision term of the Boltzmann equation is given by

$$C = \frac{\lambda^2}{2} \int \frac{d^3k_1 d^3k_2 d^3k_3}{(2\pi)^5 2^4 \omega(k_1) \omega(k_2) \omega(k_3) \omega(p)} \delta(\omega(p) + \omega(k_1) - \omega(k_2) - \omega(k_3)) \delta(\mathbf{p} + \mathbf{k}_1 - \mathbf{k}_2 - \mathbf{k}_3) \left\{ F(k_1)F(k_2)F(k_3) + F(p)F(k_2)F(k_3) - F(p)F(k_1)F(k_2) - F(p)F(k_1)F(k_3) \right\} \quad (53)$$

where we used $F(p) = f(p) + 1/2$ to simplify and combine the terms in Ref.[5].

As shown in Section III, quantum and classical propagators have different forms. The above expression (53) uses the quantum form of propagators but otherwise ignores quantum corrections. If we are in a purely classical regime, F in the above expression must be changed to f . The resulting Boltzmann equation then contains only the terms cubic in f . This is appropriate since it is easy to verify that the equilibrium distribution resulting from having only the cubic terms is $f(E) = T/E$. This form leads to the Rayleigh-Jeans catastrophe which Planck cured by inventing Quantum Mechanics.

On the other hand, if we keep the above form, the steady state solution is

$$f(E) = \frac{T}{E} - \frac{1}{2} \quad (54)$$

which may be recognized as the first 2 terms of the expansion of the Bose-Einstein distribution in the large T limit. Therefore in the $T \gg E$ limit, this form is a better approximation of the true quantum distribution than the Rayleigh-Jeans form. However unlike the Rayleigh-Jeans form, this form of the distribution function becomes negative when $T < 2E$ which is clearly unphysical. Therefore mixing the quantum propagators and the classical vertices does not necessarily improve the purely classical result.

Let us now consider why keeping the $1/2$ term in the propagator somewhat mysteriously reproduce the quantum statistical result up to the next-to-leading order terms in f . Suppose we have a scattering process that involves m initial particles (k_i) and n final particles (p_j). The quantum Boltzmann equation must contain the following combination of the density functions:

$$C_Q = \prod_{i=1}^m f_Q(k_i) \prod_{j=1}^n [1 + f_Q(p_j)] - \prod_{j=1}^n f_Q(p_j) \prod_{i=1}^m [1 + f_Q(k_i)] \quad (55)$$

to be consistent with the equilibrium Bose-Einstein density. Expanding in terms of f_Q 's (we are in the large f_Q limit), we have

$$\begin{aligned} C_Q = & \sum_{s=1}^n \prod_{i=1}^m f_Q(k_i) \prod_{j=1, j \neq s}^n f_Q(p_j) + \sum_{s=1}^n \sum_{t=1}^n \theta(s > t) \prod_{i=1}^m f_Q(k_i) \prod_{j=1, j \neq s, j \neq t}^n f_Q(p_j) \\ & - \sum_{s=1}^m \prod_{i=1, i \neq s}^m f_Q(k_i) \prod_{j=1}^n f_Q(p_j) - \sum_{s=1}^m \sum_{t=1}^m \theta(s > t) \prod_{i=1, i \neq s, i \neq t}^m f_Q(k_i) \prod_{j=1}^n f_Q(p_j) \\ & + o(f^{m+n-3}) \end{aligned} \quad (56)$$

where we defined $\theta(\text{condition}) = 1$ if the condition is met and $\theta(\text{condition}) = 0$ otherwise.

On the other hand, the classical counter-part must contain

$$C_{\text{cl}} = \sum_{s=1}^n \prod_{i=1}^m f_{\text{cl}}(k_i) \prod_{j=1, j \neq s}^n f_{\text{cl}}(p_j) - \sum_{s=1}^m \prod_{i=1, i \neq s}^m f_{\text{cl}}(k_i) \prod_{j=1}^n f_{\text{cl}}(p_j) \quad (57)$$

to be consistent with the Rayleigh-Jeans form of the equilibrium distribution. It is clear that the leading order terms match between the quantum and the classical cases. Now let $f_{\text{cl}} \rightarrow f_{\text{cl}} + 1/2$ to get

$$C'_{\text{cl}} = \sum_{s=1}^n \prod_{i=1}^m [f_{\text{cl}}(k_i) + 1/2] \prod_{j=1, j \neq s}^n [f_{\text{cl}}(p_j) + 1/2] - \sum_{s=1}^m \prod_{i=1, i \neq s}^m [f_{\text{cl}}(k_i) + 1/2] \prod_{j=1}^n [f_{\text{cl}}(p_j) + 1/2] \quad (58)$$

Expanding up to the next order in f_{cl} yields

$$C'_{\text{cl}} = \sum_{s=1}^n \prod_{i=1}^m f_{\text{cl}}(k_i) \prod_{j=1, j \neq s}^n f_{\text{cl}}(p_j) - \sum_{s=1}^m \prod_{i=1, i \neq s}^m f_{\text{cl}}(k_i) \prod_{j=1}^n f_{\text{cl}}(p_j)$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{s=1}^n \sum_{t=1}^n \theta(s \neq t) \prod_{i=1}^m f_{\text{cl}}(k_i) \prod_{j=1, j \neq s, j \neq t}^n f_{\text{cl}}(p_j) - \frac{1}{2} \sum_{s=1}^m \sum_{t=1}^m \theta(s \neq t) \prod_{i=1, i \neq s, i \neq t}^m f_{\text{cl}}(k_i) \prod_{j=1}^n f_{\text{cl}}(p_j) \\
& + o(f^{m+n-3})
\end{aligned} \tag{59}$$

which can be recognized to be the same as Eq.(56) once we use the fact that $\theta(s > t)$ is equivalent to $\theta(s \neq t)/2$ under the sums. Thus the *prescription*, $f_{\text{cl}} \rightarrow f_{\text{cl}} + 1/2$ does produce the first two terms in the quantum Boltzmann equation for any general scattering process. This also explains why the same kind of consideration also seems to work for hot QCD [17] Although it is curious that this prescription can indeed get the next-to-leading order term in f right, this is irrelevant in getting the right equilibrium limit since that requires retaining all the terms in f .

V. SEPARATION OF HARD AND SOFT MODES

In view of the discussion in the last section, the better application of the statistical mechanics of the classical field is not to get the final equilibrium distribution¹ but to use it as an intermediate stage where the field degrees of freedom and the particle degrees of freedom intermix. In this section, we illustrate how this may be achieved. A more detailed analysis of this important problem, however, is out of the scope of the present manuscript and will be reported in later publications[18].

As shown in the last section, the quantum perturbative correction becomes the same size as the classical perturbative correction when $f \sim 1$. As indicated by the equilibrium form of f , $f(k)$ becomes small as k becomes larger. Therefore, the classical field description is usually appropriate only for the soft modes.

With the price of introducing a cutoff Λ (where f becomes ~ 1), the quantum field should then be split into the soft and hard modes:

$$\varphi = \varphi_s + \varphi_h \quad \text{and} \quad \chi = \chi_s + \chi_h \tag{60}$$

where the soft modes (with the subscript s) have $|\mathbf{k}| < \Lambda$ and the hard modes (with the subscript h) have $|\mathbf{k}| \geq \Lambda$. The soft modes should then be treated as a classical field and the hard modes should be kept as a quantum field.

¹ This was also recognized in Ref.[5] where it was implied that their formalism is for the system evolving toward equilibrium but not applicable near the kinetic equilibrium.

To be concrete, consider the $\lambda\phi^4$ theory with the Hamiltonian given by

$$H = \int d^3x \left(\frac{\pi^2}{2} + \frac{(\nabla\phi)^2}{2} + \frac{m^2}{2}\phi^2 + \frac{\lambda}{4!}\phi^4 \right) \quad (61)$$

The continuum version of Eq.(23) is

$$\begin{aligned} \rho_W[\varphi_{N+1}, \pi_{N+1}^\varphi] &= \int [d\varphi_0][d\chi_0] \rho_W[\varphi_0, \pi_0^\varphi] \\ &\times \int \mathcal{D}\varphi \int \mathcal{D}\chi \int \mathcal{D}\pi^\varphi \int \mathcal{D}\pi^\chi \\ &\times \exp \left[i \int \pi^\chi (\dot{\varphi} - \pi^\varphi) - i \int \left\{ \chi \left(\dot{\pi}^\varphi - \nabla^2\varphi + m^2\varphi + \frac{\lambda}{3!}\varphi^3 \right) + \frac{\lambda}{4!}\chi^3\varphi \right\} \right] \end{aligned} \quad (62)$$

Separating the hard and the soft modes, the interaction term becomes

$$V = \frac{\lambda}{3!}(\chi_h + \chi_s)(\varphi_h + \varphi_s)^3 + \frac{\lambda}{4!}(\varphi_h + \varphi_s)(\chi_h + \chi_s)^3 \quad (63)$$

Keeping only the terms that contain the most number of φ_s yields

$$V = \frac{\lambda}{4!} \left(4\chi_s\varphi_s^3 + 4\chi_h\varphi_s^3 \right) \quad (64)$$

These terms are linear in χ_s and χ_h . Hence, the equations of motion are

$$\partial^2\varphi_s - m^2\varphi_s - \frac{\lambda}{3!}\varphi_s^3 = 0 \quad (65)$$

and

$$\partial^2\varphi_h - m^2\varphi_h = \frac{\lambda}{3!}\varphi_s^3 \quad (66)$$

These equations describe a system where the classical field evolves independent of the particles but the *quantum mechanical* free particles are produced by a classical source. From the analogous problem in QED (generation of photons from a classical source), it is an easy matter to show that the spectrum of hard modes at the end of evolution is

$$f_h(E_k, \mathbf{k}_h) = \frac{\lambda^2}{36} |J(E_k, \mathbf{k}_h)|^2 \quad (67)$$

where

$$J(E_k, \mathbf{k}_h) = \int d^3x \int_{t_0}^{\infty} dt e^{-i\mathbf{k}_h \cdot \mathbf{x} + iE_k t} (\varphi_s(t, x))^3 \quad (68)$$

This is, of course, in addition to the hard modes which already existed in the initial state.

If the cut-off Λ remains constant at all times, this is the complete solution for the hard spectrum provided that we can solve the soft mode classical equation of motion either perturbatively or non-perturbatively. In reality, it is not as simple due to the fact that the cut-off Λ should be a functional of φ_s , too. In the Color Glass Condensate approach to the heavy ion collisions [19], a similar conversion of field degrees of freedom to particle degrees of freedom was performed to get partons out of classical non-Abelian gauge field. The line of argument given here may provide a firmer ground for such a treatment. One can conceive that a kind of ‘renormalization group’ equation in the manner of Ref.[20] should exist for the cut-off. This will then enable us to describe the whole system in a consistent fashion. This topic is currently under investigation.

VI. SUMMARY AND DISCUSSION

In this paper, we showed that one can indeed go from the statistical mechanics of quantum field theory to the statistical mechanics of particles via classical field theory. Along the way, we encountered several subtleties that had to be carefully dealt with. In this regard, we improve upon the treatment given in Ref.[5].

Going from quantum field theory to the classical field theory involves the Wigner functional which plays the role of initial density function. In other words, the probability of an initial state depends on both the value of the field and the value of the first time derivative. The form of the initial Wigner functional in turn determines the propagators. In addition, our ability to use the Feynman diagrams with energy-momentum conservation at each vertex depends very much on the fact that time translation invariance is maintained. For this to work at least approximately, the Wigner transformed propagators $G_W(p, X)$ (*c.f.* Eq.(37)) cannot depend strongly on $X = (x + y)/2$. This in turn implies that the density matrix at time t must also be approximately of the form

$$\ln \rho \sim E_k^2 \varphi_k^2 + \pi_k^2 \sim H_k \quad (69)$$

where φ_k and π_k are Fourier components of the corresponding fields at t and H_k is the corresponding free field Hamiltonian. This is a rather strong condition which is in general valid only near equilibrium. Whether this can be reconciled with the classical field limit is not yet clear and requires further study.

Going from the quantum field theory to the classical field theory also involves a Jacobian in the form of the Van Vleck determinant. The role of this Jacobian turned out to be to cancel the single remaining quantum effect still left in the classical perturbation theory in the form of the tadpole self-energy.

It must be also emphasized that to define propagators, one must define what is meant by the ‘un-perturbed vacuum’. Pure classical vacuum cannot fluctuate as it is literally a state where nothing exists. On the other hand, the quantum vacuum fluctuates all the time. The importance of this distinction is particularly apparent when one considers the free field theory. The form of the path integral for the classical and the quantum theory in this case is exactly the same except the form of the initial density functional. To be consistent, the classical initial density functional must conform to the classical vacuum property and the quantum density functional must conform to the quantum vacuum property. Mixing of the two formalism leads to an interesting conclusion that the classical field theory can reproduce the results of *quantum statistical* Boltzmann equation up to the next leading order in the density (Ref.[5]). The correct Boltzmann equation obtained from classical perturbation theory of course yields the Rayleigh-Jeans form of the equilibrium distribution function as it must.

As for the validity of the classical perturbation theory, in addition to the two conditions already mentioned we also argued that there is a limit that a purely classical perturbation theory makes sense. For instance, we have shown that if $f \sim 1/\sqrt{\lambda}$, the leading order quantum correction is as big as the second order classical correction. And hence, going beyond the first order classical perturbation theory doesn’t make much sense. Quantum effect must be considered after the leading order.

We would also like comment on the meaning of ‘thermalization’ here. It is well known that an isolated system *as a whole* cannot thermalize. A simple example is an eigenstate of the total Hamiltonian which, by definition, is stationary. This is in contrast to the Boltzmann equation where the stationary solution is guaranteed to be the equilibrium distribution. The crucial difference of course is whether one is interested in the system as a whole, or just a *part* of the system be it in the momentum space or the coordinate space. It is therefore perhaps more natural to apply the current formalism to a particular sector of the system,

say the soft modes, and regard the hard spectrum as particle degrees of freedom².

In summary, in this paper, we re-examined the derivation of the Boltzmann equation from the classical field theory as advocated by Mueller and Son [5]. We pointed out a few subtle points in manipulating functional integrals and developing the perturbation theory and showed how to deal with them. With our improvement, the framework advocated by Mueller and Son has a potential to be a very useful tool in investigating dense many-body systems, for instance, the problem of converting field degrees of freedom to particle degrees of freedom. Further study in this line is continuing and will be reported in later publications.

Acknowledgments

The author gratefully acknowledges C. Gale and G. Moore for many discussions. He also thanks R. Venugopalan for his helpful suggestions in preparing this manuscript. The author also thanks D. Kharzeev and G. Aarts for their suggestions. S.J. is supported in part by the Natural Sciences and Engineering Research Council of Canada and by le Fonds Nature et Technologies of Québec. S.J. also thanks RIKEN BNL Center and U.S. Department of Energy [DE-AC02-98CH10886] for providing facilities essential for the completion of this work.

-
- [1] L. D. McLerran and R. Venugopalan, Phys. Rev. D **49**, 2233 (1994); *ibid.*, 3352, (1994); *ibid.*, **50**, 2225 (1994); *ibid.*, **59**:094002, (1999).
 - [2] A. Kovner, L. D. McLerran and H. Weigert, Phys. Rev. D **52**, 3809 (1995)
 - [3] A. Kovner, L. D. McLerran and H. Weigert, Phys. Rev. D **52**, 6231 (1995)
 - [4] J. P. Blaizot and E. Iancu, Nucl. Phys. B **557**, 183 (1999) [arXiv:hep-ph/9903389].
 - [5] A. H. Mueller and D. T. Son, Phys. Lett. B **582**, 279 (2004)
 - [6] E. Calzetta and B. L. Hu, Phys. Rev. D **37**, 2878 (1988).
 - [7] S. Mrowczynski and B. Muller, Phys. Rev. D **50**, 7542 (1994)
 - [8] E. Deotto, E. Gozzi and D. Mauro, J. Math. Phys. **44**, 5902 (2003)

² This is different from similar separation of hard and soft modes previously considered in the *equilibrium setting*. For instance, see Ref.[21].

- [9] E. Deotto, E. Gozzi and D. Mauro, J. Math. Phys. **44**, 5937 (2003)
- [10] A.N. Drozdov Phys. Rev. E **55**, 2496 (1997).
- [11] E. Gozzi and M. Regini, Phys. Rev. D **62**, 067702 (2000).
- [12] G. Aarts and J. Smit, Nucl. Phys. B **511**, 451 (1998)
- [13] G. Aarts, B. J. Nauta and C. G. van Weert, Phys. Rev. D **61**, 105002 (2000)
- [14] J. Berges, Nucl. Phys. A **702**, 351 (2002)
- [15] J. Berges, AIP Conf. Proc. **739**, 3 (2005)
- [16] R.P. Feynman and A.R. Hibbs, *Quantum mechanics and path integrals*, 1965, McGraw-Hill
- [17] T. Stockamp, *Classical approximation of the Boltzmann equation in high energy QCD*, arXiv:hep-ph/0408206.
- [18] Steffen Weinstock, Raju Venugopalan and Sangyong Jeon in preparation.
- [19] A. Krasnitz and R. Venugopalan, Phys. Rev. Lett. **86**, 1717 (2001)
- [20] D. Boyanovsky, H. J. de Vega, R. Holman and M. Simionato, Phys. Rev. D **60**, 065003 (1999)
- [21] M. Gleiser and R. O. Ramos, Phys. Rev. D **50**, 2441 (1994)